



# REVISIONI SISTEMATICH E META-ANALISI

Coordinatore:  
**Dr.ssa Stefania Gori**

*Evento ECM MODULO 3  
(formazione di base)*



**NEGRAR**  
**6-7 Maggio 2016**  
Centro Formazione  
Ospedale Sacro Cuore  
Don Calabria

# SOF: Summary of Findings tables

Negrar, 06 maggio 2016

**Formulate question**


P  
I  
C  
O

Select outcomes  
Rate importance

Outcome Critical  
Outcome Critical  
Outcome Important  
Outcome Not important

1

**Outcomes across studies**



2

**Create evidence profile with GDT**

Quality assessment	Summary of findings																
<table border="1"> <tr> <th>High</th> <th>Moderate</th> <th>Low</th> <th>Very low</th> </tr> <tr> <td>100%</td> <td>75%</td> <td>50%</td> <td>25%</td> </tr> </table>	High	Moderate	Low	Very low	100%	75%	50%	25%	<table border="1"> <tr> <th>High</th> <th>Moderate</th> <th>Low</th> <th>Very low</th> </tr> <tr> <td>100%</td> <td>75%</td> <td>50%</td> <td>25%</td> </tr> </table>	High	Moderate	Low	Very low	100%	75%	50%	25%
High	Moderate	Low	Very low														
100%	75%	50%	25%														
High	Moderate	Low	Very low														
100%	75%	50%	25%														

Summary of findings & estimate of effect for each outcome

3

**Rate quality of evidence for each outcome**

High  
Moderate  
Low  
Very low

Randomization raises initial quality  
RCTs: high  
Observational: low

Grade down


1. Risk of bias
2. Inconsistency
3. Indirectness
4. Imprecision
5. Publication bias

Grade up

1. Large effect
2. Dose response
3. Opposing bias & Confounders

4

**Grade overall quality of evidence across outcomes based on lowest quality of critical outcomes**



6

**Guideline/Decision**

**Evidence synthesis (SR, HTA)**

**Recommendation/Decision**

**Grade recommendations (Evidence to Decision)**

- For or against (direction) ↓↑
- Strong or conditional/weak (strength)

By considering balance of consequences (evidence to recommendations):

5

- Quality of evidence
- Balance benefits/harms
- Values and preferences
- Feasibility, equity and acceptability
- Resource use (if applicable)

**EtD framework with GRADEpro**

Issue	Assessment	Research evidence	Explanation/Justification/Definition
What is the overall quality of evidence?	High, Moderate, Low, Very low	100% High, 75% Moderate, 50% Low, 25% Very low	...



**Formulate Recommendations/Decision**

- "The panel recommends that ....should..."
- "The panel suggests that ....should..."
- "The panel suggests to not ..."
- "The panel recommends to not..."

**Transparency, clear, actionable Research?**

# What are we grading?

- two components
- quality of body of evidence
  - extent to which confidence in estimate of effect adequate to support decision
    - high, moderate, low, very low
- strength of recommendation
  - strong and weak

# RISK OF BIAS/QUALITY OF EVIDENCE

- **RISK OF BIAS:** concerns with the internal validity of study results, i.e. the systematic (as opposed to random) deviation of the results of a study from the 'true' results, which is caused by the way the study is designed or conducted

# Risk of bias table for RCTs Cochrane Collaboration

Bias	Authors' judgement	Support for judgement
<b>Random sequence generation (selection bias)</b>	Low risk of bias	
	High risk of bias	
	Unclear risk of bias	
<b>Allocation concealment (selection bias)</b>	Low risk of bias	
	High risk of bias	
	Unclear risk of bias	
<b>Blinding of participants and personnel (performance bias)</b>	Low risk of bias	
	High risk of bias	
	Unclear risk of bias	
<b>Blinding of outcome assessment (detection bias)</b>	Low risk of bias	
	High risk of bias	
	Unclear risk of bias	
<b>Incomplete outcome data (attrition bias)</b>	Low risk of bias	
	High risk of bias	
	Unclear risk of bias	
<b>Selective reporting (reporting bias)</b>	Low risk of bias	
	High risk of bias	
	<b>Unclear risk of bias</b>	

# RISK OF BIAS/QUALITY OF EVIDENCE

- **QUALITY OF EVIDENCE** (according to GRADE): more extensive evaluation considering also other domains :
- **risk of bias**
- inconsistency of the results across the studies
- imprecision
- indirectness
- risk of publication bias
- magnitude of the effect
- dose response gradient
- Residual confounding

# Quality of evidence

- GRADE is “**outcome centric**”:
  - rating is made for each outcome, and quality **may differ** -indeed, is likely to differ - **from one outcome to another within a single study and across a body of evidence**
- E.g: **subjective** outcomes are prone to performance and detection bias, while **objective** outcomes are not
- E.g. **one outcome** within a review **could have imprecision** in the pooled estimate of the effect, while **another could have not**

# grades of evidence and Interpretation

Symbol	Quality	Interpretation
⊕⊕⊕⊕	High	We are very confident that the true effect lies close to that of the estimate of the effect
⊕⊕⊕○	Moderate	We are moderately confident in the effect estimate: The true effect is likely to be close to the estimate of the effect, but there is a possibility that it is substantially different
⊕⊕○○	Low	Our confidence in the effect estimate is limited: The true effect may be substantially different from the estimate of the effect
⊕○○○	Very low	We have very little confidence in the effect estimate: The true effect is likely to be substantially different from the estimate of effect



# Evidence profile and Summary of findings

**Evidence profile:** summary of evidence for a given question with a detailed quality assessment and an explicit judgment of each factor that determines the quality. Used by guideline producers

**Summary of findings:** summary of evidence for a given question with quality assessment but not the detailed judgments. Prepared within SRs

## For each outcome:

- a) Relative risk with 95%CI
- b) Absolute risk : **SoF** presents the **absolute risks in intervention and control groups** with a CI around the intervention group rate, **EP** presents the **risk difference** with 95%CI.
- c) Number of participants (n of studies) included

# Evidence profile: use of antibiotics (penicillin) versus no use of antibiotics in children with sickle cell disease. Source: Hirst et al. 4

Quality assessment							No of patients		Effect		Quality	Importance
No of studies	Study design	Risk of bias	Inconsistency	Indirectness	Imprecision	Other considerations	Penicillin prophylaxis	Standard care	Relative (95% CI)	Absolute (95% CI)		
Incidence of pneumococcal infection, for initiation of treatment - Initiation of penicillin												
2	Randomized trials	Not serious <sup>1</sup>	serious <sup>2</sup>	not serious <sup>3</sup>	serious <sup>4</sup>	none <sup>5</sup>	9/248 (3.6%)	19/209 (9.1%)	<b>OR 0.37</b> (0.16 to 0.86)	55 fewer per 1000 (from 12 fewer to 75 fewer)	⊕⊕○○ LOW	CRITICAL
Deaths, for initiation of treatment - Initiation of penicillin												
1	randomized trials	not serious <sup>6</sup>	not serious	not serious	serious <sup>4</sup>	none <sup>5</sup>	0/105 (0.0%)	4/110 (3.6%)	<b>OR 0.11</b> (0.01 to 2.11)	32 fewer per 1000 (from 36 fewer to 37 more)	⊕⊕⊕○ MODERATE	CRITICAL
Adverse drug effects - Nausea and vomiting												
1	randomized trials	not serious <sup>6</sup>	not serious	not serious	serious <sup>4</sup>	none <sup>5</sup>	2/201 (1.0%)	1/199 (0.5%)	<b>OR 1.99</b> (0.18 to 22.12)	5 more per 1000 (from 4 fewer to 95 more)	⊕⊕⊕○ MODERATE	CRITICAL

1. blinding and concealment were not clear for one of the two studies
2. heterogeneity exists; p-value for testing heterogeneity is 0.07 and I<sup>2</sup>=69%
3. the question addressed is the same for the evidence regarding the population, intervention, comparator and outcome
4. total sample size is small and the total number of events is <300 (a threshold rule-of-thumb value)
5. insufficient number of studies to assess publication bias
6. unclear allocation concealment

# Summary of findings

Summary of finding: antibiotics for acute otitis media in children

Antibiotics compared with placebo for acute otitis media in children

Patient or population: Children with acute otitis media

Setting: High- and middle-income countries


Intervention: Antibiotics

Comparison: Placebo

Outcomes	Estimated risks (95% CI)		Relative effect (95% CI)	No. of Participants (studies)	Quality of the evidence (GRADE)
	Control risk <sup>a</sup> Placebo	Intervention risk Antibiotics			
Pain at 24h	367 per 1,000	330 per 1,000 (286–382)	RR 0.9 (0.78–1.04)	1229 (5)	⊕⊕⊕⊕ High
Pain at 2–7 d	257 per 1,000	185 per 1,000 (159–213)	RR 0.72 (0.62–0.83)	2791 (10)	⊕⊕⊕⊕ High
Hearing, inferred from the surrogate outcome abnormal tympanometry—1 mo	350 per 1,000	311 per 1,000 (262–375)	RR 0.89 (0.75–1.07)	927 (4)	⊕⊕⊕○ Moderate <sup>b</sup>
Hearing, inferred from the surrogate outcome abnormal tympanometry—3 mo	234 per 1,000	227 per 1,000 (178–290)	RR 0.97 (0.76–1.24)	808 (3)	⊕⊕⊕○ Moderate <sup>b</sup>
Vomiting, diarrhea, or rash	113 per 1,000	156 per 1,000 (123–199)	RR 1.38 (1.09–1.76)	1,401 (5)	⊕⊕⊕○ Moderate <sup>c</sup>

# Rating quality of evidence

GRADE's approach begins with the study design. Randomized controlled trials (RCTs) start as high-quality evidence and observational studies as low-quality evidence supporting estimates of intervention effects.



# Determinants of quality/certainty of a body of evidence

- RCTs ⊕⊕⊕⊕


- observational studies ⊕⊕○○

- **5 factors that can lower quality**

1. limitations in detailed study design and execution (*risk of bias criteria*)
2. Inconsistency (*or heterogeneity*)
3. Indirectness (*PICO and applicability*)
4. Imprecision
5. Publication bias

- **3 factors can increase quality**

1. large magnitude of effect
2. opposing plausible residual bias or confounding
3. dose-response gradient



# 1. risk of bias

Deferasirox for managing transfusional iron overload in people with sickle cell disease (Review)

	Random sequence generation (selection bias)	Allocation concealment (selection bias)	Blinding (performance bias and detection bias)	Incomplete outcome data (attrition bias)	Selective reporting (reporting bias)	Other bias
Vichinsky 2007	?	+	-	-	-	?
Vichinsky 2011	?	?	-	-	+	?

# Risk of bias

- Outcome specific
  - **each trial contributes** toward the estimate of magnitude of effect.
  - **larger trials with many events** will contribute **more** ( look at the weight of each study in the forest plot)

## 2. Inconsistency (heterogeneity) between studies results


- Variation in size of effect ( **Point estimates vary widely** across studies)
- **Confidence intervals** (CIs) show minimal or **no overlap**
- The statistical test for heterogeneity which tests the null hypothesis that all studies in a meta-analysis have the same underlying magnitude of effect shows a low **P-value (< 0.05)**
- The **I<sup>2</sup>** which quantifies the proportion of the variation in point estimates due to among-study differences (< 40% : low, 30 e 60% : moderate, **50%** e 90% : substantial, 75 e 100% : considerable)
- **All statistical approaches have limitations**, and their **results should be seen in the context of a subjective** examination of the variability in point estimates and the overlap in CIs.



Study or Subgroup	Any antipsychotic			Placebo			Weight	Mean Difference
	Mean	SD	Total	Mean	SD	Total		IV, Random, 95% CI
Kampman 2003	1.3	0.8	15	1	0.9	15	28.3%	0.30 [-0.31, 0.91]
Reid 2005	6.1	2.1	16	3.5	2.3	15	20.3%	2.60 [1.05, 4.15]
Tapp 2015	0.4	1.26	29	1.63	2.39	31	25.6%	-1.23 [-2.19, -0.27]
Winhusen 2007	2.8	2.45	60	3.44	2.76	59	25.8%	-0.64 [-1.58, 0.30]
<b>Total (95% CI)</b>			<b>120</b>			<b>120</b>	<b>100.0%</b>	<b>0.13 [-1.08, 1.35]</b>

Heterogeneity:  $\tau^2 = 1.26$ ;  $\text{Chi}^2 = 19.78$ ,  $df = 3$  ( $P = 0.0002$ );  $I^2 = 85\%$

Test for overall effect:  $Z = 0.22$  ( $P = 0.83$ )



# 3. Directness of Evidence

## generalizability, transferability, applicability


- differences between PICO and available evidence in
  - populations/patients (interested in children found adults population)
  - interventions (interested in high dosage, found low dosage, interested in long treatment, found short, etc)
  - outcomes (interested in important but we found surrogate; e.g hip fracture vs bone density; interested in long term but found short term results)
- indirect comparisons
  - interested in A versus B
  - found A versus C and B versus C

# 4. Publication Bias

Consider rating down if:

- You find **systematic reviews performed early**, when only few initial studies are available, that will overestimate effects when “negative” studies face delayed publication. Early positive studies, particularly if small in size, are suspect.
- You find **only small “positive” studies**, mainly if sponsored by industry
- **Funnel plot showing asymmetry** but
- Funnel plot should be seen as a generic means of displaying small-study effects – a tendency for the intervention effects estimated in smaller studies to differ from those estimated in larger studies (Sterne 2000). Small-study effects may be due to reasons other than publication bias ( low methodological quality, chance, patients characteristics).
- **Funnel plot should be used only when there are at least 10 studies** included in the meta-analysis, because when there are fewer studies the power of the tests is too low to distinguish chance from real asymmetry


# Funnel plot



# Funnel plot

- On the **horizontal axis** : measure of treatment effect
- On the **vertical axis**: standard error ( SE) of the intervention effect estimate: measure of **precision of the estimate** ; SE is determined by sample size, and by the number of participants experiencing the event for dichotomous outcomes, and the standard deviation of responses for continuous outcomes.
- Precision of the estimated intervention effect increases as the size of the study increases. Effect estimates from small studies will therefore scatter more widely at the bottom of the graph, with the spread narrowing among larger studies. **In the absence of bias the plot should approximately resemble a symmetrical funnel plot.**

# Funnel plot



## 5. Imprecision of the overall estimate

- **Wide confidence intervals** (CIs inform the impact of random error on evidence quality; CI expresses the range in which the truth plausibly lies)
- **Small number of events**
- **Small sample size**


# Optimal information size

- **We suggest the following:**
- **if the total number of patients included in a systematic review is less than the number of patients generated by a conventional sample size calculation for a single adequately powered trial, consider rating down for imprecision.** Authors have referred to this threshold as the “optimal information size” (OIS)



Required sample size (assuming  $\alpha$  of 0.05, and  $\beta$  of 0.2) for RRR of 20%, 25%, and 30% across varying control event rates. For example, **if the best estimate of control event rate was 0.2 and one specifies an RRR of 25%, the OIS is approximately 2,000 patients** (GRADE guideline n.6 Journal of Clinical Epidemiology 64 (2011))

1282017021



Optimal information size given  $\alpha$  of 0.05 and  $\beta$  of 0.2 for varying control event rates and relative risks.

# Power is more closely related to number of events than to sample size

( GRADE guideline n.6 Journal of Clinical Epidemiology 64 (2011) 1283e1293)

Calculating the OIS for **dichotomous outcome** requires specifying:

- probability of detecting a false effect – type I error ( $\alpha$ ; usually 0.05)
- probability of detecting a true effect – power (usually 80% [power = 1 – type II error;  $\beta$ ; usually 0.20])
- realistic relative risk reduction (RRR; we suggest a default of 25%)
- control event rate (we suggest the median of the available trials, or the rate from a dominating trial, if it exists).




Table 1: Optimal information size implications from Figure above

Total Number of Events	Relative Risk Reduction	Implications for meeting OIS threshold
100 or less	$\leq 30\%$	Will almost never meet threshold whatever control event rate
200	30%	Will meet threshold for control event rates for ~ 25% or greater
200	25%	Will meet threshold for control event rates for ~ 50% or greater
200	20%	Will meet threshold only for control event rates for ~ 80% or greater
300	$\geq 30\%$	Will meet threshold
300	25%	Will meet threshold for control event rates ~ 25% or greater
300	20%	Will meet threshold for control event rates ~ 60% or greater
400 or more	$\geq 25\%$	Will meet threshold for any control event rate
400 or more	20%	Will meet threshold for control event rates of ~ 40% or greater

# OIS for continuous outcomes

- Authors can calculate the OIS for continuous variables **in exactly the same way they can for binary variables by specifying the  $\alpha$  and  $\beta$  errors** (we have suggested 0.05 and 0.2) and the  $\Delta$  ( i.e. the difference one wishes to detect as clinically relevant ), and choosing an appropriate standard deviation from one of the relevant studies.
- A particular challenge in calculating the OIS for continuous variables arises when studies have used different instruments to measure a construct, and the pooled estimate is calculated using a standardized mean difference.
- we suggest authors choose one of the available instruments (ideally, one in which an estimate of the minimally important difference is available) and calculate an OIS using that instrument

# OIS for continuous outcomes

whenever there are **sample sizes** that are **less than 400**, review authors and guideline developers should certainly consider **rating down** for imprecision.

# Downgrading and OIS

- if OIS **not met** downgrade for imprecision
- If OIS **met and** the 95% CI excludes a relative risk (RR) of 1.0 (**statistically significant results**), precision is adequate.
- **if OIS met but the 95% CI includes a RR of 1 ( null effect) ,** authors should **consider whether CIs include appreciable benefit or harm** (we suggest a RR of under **0.75** or over **1.25** as a rough guide) ; if yes downgrading for imprecision may be appropriate.

# What can raise quality?

1. **large magnitude of effect** can upgrade (**RRR 50%/RR 2**)
  - **very large two levels (RRR 80%/RR 5)** ; modeling studies suggests that **confounding** (from nonrandom allocation) alone is **unlikely to explain** associations with a relative risk (RR) greater than 2 (or less than 0.5), and very unlikely to explain associations with an RR greater than 5 (or less than 0.2)
  - Es: relationship between infant **sleeping position and sudden infant death** syndrome (SIDS) found an odds ratio (OR) of 4.1 (95% confidence interval [CI]: 3.1, 5.5) of SIDS occurring with front vs. back sleeping positions

# What can raise quality?

## 2. dose response relation

- higher INR – increased bleeding
- childhood lymphoblastic leukemia
  - risk for CNS malignancies 15 years after cranial irradiation
  - no radiation: 1% (95% CI 0% to 2.1%)
  - 12 Gy: 1.6% (95% CI 0% to 3.4%)
  - 18 Gy: 3.3% (95% CI 0.9% to 5.6%)



# Residual confounding

- 3. all plausible residual confounding **may be working to reduce the demonstrated effect or increase the effect if no effect was observed** (underestimate of the treatment effect)
- Es: effect of condom use on HIV infection among men who have sex with men RR: 0.34 [0.21, 0.54] **(RRR: 66%) in favor of condom use** compared with no condom use. Condom users were more likely to have more partners **(but studies did not adjust for this confounding factor in their analyses)**. Considering the number of partners would, if anything, strengthen the effect estimate in favor of condom use.